

# <sup>1</sup>H NMR STUDY OF THE CONFORMATION OF THE RIBOSE PHOSPHATE MOIETY OF 6-AZAURIDINE-5'-MONOPHOSPHATE – A NUCLEOTIDE WITH AN UNUSUAL CONFORMATION

Frank E. HRUSKA, Donald J. WOOD, Richard J. MYNOTT and Ramaswamy H. SARMA

*Department of Chemistry, University of Manitoba, Winnipeg, Manitoba, Canada*

and

*Department of Chemistry, The State University of New York at Albany, Albany, New York 12222, USA*

Received 2 February 1973

## 1. Introduction

6-Azauridine-5'-monophosphate (5'-azaUMP, I) is known to be a potent antileukemic agent [1] and to determine whether this activity may be correlated with an unusual conformation, we have examined its molecular geometry in solution using <sup>1</sup>H NMR spectroscopy. Comparison of the <sup>1</sup>H NMR data on U (uridine), 5'-UMP, 6-azauridine (azaU) and 5'-azaUMP, show that the presence of the 6-aza base has a marked destabilizing influence upon the gauche-gauche (II) conformation constrained to the C(4') – C(5') bond of 5'-azaUMP.

Table 1

Coupling constants in Hz for U, 6-azaU, 5'-UMP and 5'-azaUMP, D<sub>2</sub>O solutions, 30°.

Nuclei	U	aza U	5'-UMP	5'-azaUMP
J <sub>1'2'</sub>	4.4	3.6	5.3	4.2
J <sub>2'3'</sub>	5.3	5.2	5.0	5.4
J <sub>3'4'</sub>	5.5	5.8	3.9	5.4
J <sub>4'5'</sub>	3.0	3.2	2.5	4.6
J <sub>4'5''</sub>	4.4	5.6	3.0	6.1
J <sub>5'5''</sub>	-12.7	-12.4	-13.5	-11.3
J <sub>1'5</sub>	0.35	0.61	<0.2	0.55
J <sub>P4'</sub>	—	—	1.8	<0.2
J <sub>P5'</sub>	—	—	3.2	5.6
J <sub>P5''</sub>	—	—	5.1	6.1
J <sub>4'5'</sub> + J <sub>4'5''</sub>	7.4	8.8	5.5	10.7
J <sub>P5'</sub> + J <sub>P5''</sub>	—	—	8.3	11.7

## 2. Materials and methods

Spectra of U, azaU, 5'-UMP and 5'-azaUMP (commercial products), 0.1 M, pD 8.0, 30°, in D<sub>2</sub>O, were obtained using either a 220 MHz, continuous wave or a 100 MHz, fast Fourier transform system. Spectra were recorded both with coupling to the <sup>31</sup>P of the phosphate and with <sup>31</sup>P decoupling and were analyzed using the computer program LAME. In fig. 1 we have illustrated the experimental and computer simulated spectrum (only the ribose region) of 5'-azaUMP. The data are compiled in the table.

## 3. Results and discussion

Population distribution of the rotamers constrained to the C(4') – C(5') bond (II–IV) can be evaluated from the experimental value of J<sub>4'5'</sub> + J<sub>4'5''</sub> (Σ) [2, 3]. Any perturbations resulting in an *increase* in the g–g rotamer populations should be manifested by a *decrease* in the experimental *sum* J<sub>4'5'</sub> + J<sub>4'5''</sub>, whereas an increase in the g–t and/or t–g populations, at the expense of the g–g conformer, should result in an *increase* in the observed sum [2, 3]. Populations distribution of the rotamers constrained to the C(5') – O(5') bond (V–VII) can be monitored in an analogous way.

The experimental sum J<sub>4'5'</sub> + J<sub>4'5''</sub> (7.4 Hz for U; 8.8 Hz for azaU), shows that the presence of the 6-azauracil base has a destabilizing influence, relative




Fig. 1. a) Experimental spectrum of the ribose region of 6-aza-5'-UMP taken in a 100 MHz NMR system, using a 16K fast Fourier transform system. b) Computer simulated spectrum of the ribose region of 6-aza-5'-UMP.

to uracil, upon the *g-g* orientations (II) about the C(4') – C(5') bond of a nucleoside. This destabilizing effect is more conspicuous for the *monophosphate derivative* for which an increase of 5.2 Hz is observed when the uracil is substituted by the 6-azabase. A simple relationship (1) between  $P_{gg}$  (the fractional population of the *g-g* rotamer) and  $\Sigma$  (the observed sum of  $J_{4'5'} + J_{4'5''}$ ) can be derived by manipulating equations 1–3 in reference 2. This equation provides

$$P_{gg} \approx \frac{12 - \Sigma}{8} \quad (1)$$

rough estimates of *g-g* populations. Substitutions of the data into (1) leads to  $P_{gg}$  values: U(60%); azaU (40%); 5'-UMP (80%); 5'-azaUMP (15%). We have derived a similar expression (2) for the C(5') – C(5') bond of the monophosphates




$$P_{g'g'} \approx \frac{24 - \Sigma}{18} \quad (2)$$

where  $P_{g'g'}$  and  $\Sigma$  are, respectively, the populations of the *g'-g'* rotamer (V) and the observed sum  $J_{P-H(5')} + J_{P-H(5'')}$ . The coupling constants used are:  $J_{PH(gauche)} \approx 3$  Hz and  $J_{PH(trans)} = 21$  Hz [4, 5]. The derived  $P_{g'g'}$  values – 5'-UMP (90%), 5'-aza-

UMP (70%) – suggest a slight destabilizing influence of the 6-aza base upon the *g'g'* rotamer of a 5'-nucleotide.

In 5'-UMP, but *not* in 5'-azaUMP, we have observed a four bond  $J_{P4'}$ , long-range coupling. Four bond H–H coupling constants are known to be strongly dependent upon the stereochemical arrangement of the intervening bonds and largest for the all *trans* (W) conformation [6]. Hall and coworkers [4, 7] have presented clear evidence that H–P coupling in H–C–O–P fragments do have a similar stereorelationship. The “W” relationship between the phosphorus and 4' hydrogen of a 5' mononucleotide is encountered when the molecule is simultaneously oriented *g-g* and *g'-g'* (VIII), but is destroyed by rotations about either of the C(4') – C(5') or C(5') – O(5') bonds into one of the *trans* conformations. We believe that the observed  $^4J_{P4'}$  values for 5'-UMP (1.8 Hz) and 5'-azaUMP ( $\approx 0.1$  Hz) provide further evidence for the suggested destabilizing influence of a 6-azauracil base upon the *g-g* and *g'-g'* conformer of a 5' nucleotide.

The limitations of the applications of Karplus equation to decipher ribose conformation from  $J_{1'2'}$ ,  $J_{2'3'}$  and  $J_{3'4'}$  have been discussed elsewhere [8] and it has been suggested that the coupling constant data are best treated in terms of two favored modes viz C(2')



*endo* and C(3') *endo*. The J values in the table indicate that the ribofuranose ring in the nucleosides and nucleotides is not conformationally pure and it may exist as an equilibrium mixture of C(2') *endo* and C(3') *endo* modes undergoing interconversion via pseudorotation [9]. If the small differences in the magnitudes of the J values is anything of value, one may indicate that 5'-UMP displays a slight preference for the C(2') *endo* mode and 5'-azaUMP, a slight preference for the C(3') *endo* mode.

Comparison of the X-ray data, obtained by Saenger and Suck [10] with the present NMR data, show that 5-azaUMP retains its unusual solution conformation, in the solid state as well, and this is the first time one has encountered a 5' nucleotide whose exocyclic  $\text{CH}_2\text{OPO}_3^{2-}$  side chain does not follow the general

pattern [9a] of g-g, g'-g' conformation; and it is likely that the antileukemic property of 5'-azaUMP is related to this unusual conformation.

## Acknowledgements

This research was supported by grants from the National Cancer Institute of National Institutes of Health, USA (CA12462-02), the National Science Foundation (GB28015, GP28061), and the National Research Council of Canada (CI544, A6434) and the Research Corporation of New York. The authors thank Drs. B. Pullman, W. Saenger and M. Sundaralingam for illuminating discussions. We also are particularly grateful to Dr. W. Saenger for privately communicating his X-ray data by way of sending us a preprint.

## References

- [1] R.E. Hanschumacher, P. Calabresi, A.D. Welch, V. Bono, H. Fallou and E. Frei, Cancer Chemotherapy Reports 21 (1962) 1; J. Skoda, Prog. Nuc. Acid Res. 2 (1963) 197.
- [2] F.E. Hruska, A.A. Grey and I.C.P. Smith, J. Am. Chem. Soc. 92 (1970) 4088.
- [3] F.E. Hruska, A.A. Smith and J.C. Dalton, J. Am. Chem. Soc. 93 (1971) 4334.
- [4] L.D. Hall and R.B. Malcolm, Chem. Inc. 92 (1968).
- [5] M. Tsuboi, S. Takahashi, Y. Kyogoku, H. Hayatsu, T. Ukita and M. Kainosho, Science 166 (1969) 1504.
- [6] A. Rassat, C.W. Jefford, J.M. Lehn and B. Waegell, Tetrahedron Letters 233 (1964); M. Barfield and B. Chakrabarti, Chem. Rev. 69 (1969) 757.
- [7] L.D. Hall and R.B. Malcolm, Can. J. Chem. 50 (1972) 2092; 50 (1972) 2102; B. Donaldson and L.D. Hall, Can. J. Chem. 50 (1972) 2111.
- [8] R.H. Sarma and R.J. Mynott, J. Am. Chem. Soc. in press.
- [9] M. Sandaralingam, Proceedings, International Symposium on the Conformations of Biological Molecules and Polymers - Symposia on Quantum Chemistry and Biochemistry, eds: B. Pullman and E.D. Bergman, Vol. V (Academic Press) in press; L.D. Hall, Pr. R. Steiner and C. Pedersen, Can. J. Chem. 48 (1970) 1155.
- [10] W. Saenger and D. Suck, private communications, manuscript submitted for publication.